Table 4.13 Atmospheric inputs from UK sources to the North Sea[1] : 1987 - 2000

United Kingdom tonnes

Element		1987	1988	1989	1990	1991	1992	1993	1994	1995	1995	1996	1997	1998	1999	2000
Arsenic	As			210	200	140	57	57	69	57	56 r	..	70	34	38	48
Cadmium	Cd	98	100	77	67	59	31	34	29	27	23 r	..	28	12	13	19
Chromium	Cr	280	320	260	150	130	180	48	59	140	130 r	..	71	99	67	55
Copper	Cu	970	930	1,100	590	1,400	350	330	400	480	380 r	..	450	240	280	330
Nickel	Ni	360	340	440	340	450	250	150	220	290	250 r	..	67	79	72	83
Lead	Pb	3,500	3,500	3,000	1,400	2,400	880	760	980	940	880 r	..	810	430	400	392
Titanium	Ti	770	970	1,100	500	800	630	240	390	320	310 r	..	480	280	180	83
Zinc	Zn	5,400	5,300	6,100	4,900	5,800	2,800	2,700	5,700	3,100	4,100 r	..	3,300	2,300	1,400	3,500

Source: AEA Technology

[1] By method of Cambray et al, 1975. Deposition weighted average based on measured concentrations and rainfall.

Source publication: Digest of Environmental Statistics, Published March 2002
Department for Environment, Food and Rural Affairs
http://www.defra.gov.uk/environment/statistics/des/index.htm