APPENDIX VI

Overview of cetacean by-catches, fishing gears and area (reproduced from CETASEL report, De Haan et al. 1997)

| Species | Number of | Area | Fishing Method | Period | Reference | Comment |
|---------------------|---------------|-----------------------|------------------|--------------|---|--------------|
| | by-catches | | | | | |
| Killer whale | 1 | Mediterranean, Sicily | Tuna trap | 1989 | Di Natale, 1989 in Northridge, 1991 | |
| | 8 | NE Pacific | Trawl | 1973 - 1993 | Teshima & Ohsumi, 1983; Perez & Loughlin, 1991; NMFS, 1995; Matkin & | |
| | | | | | Saulitis, 1994; in Fertl & Leatherwood, 1995 | |
| | Small numbers | Alaska | Trawl | 1990 | Barlow et al., 1990, in Northridge 1991 | |
| | 1 | SW Pacific | Trawl | 1979 | Anonymous 1981, in Fertl & Leatherwood, 1995 | |
| Pilot whale | 297 | East US | Squid and | 1977 - 1988 | Waring et al., 1990 | 55% of total |
| | | | mackerel trawls | | | by-catch1 |
| | 405 | NE Atlantic | Trawl | 1977 – 1988 | Fertl & Leatherwood, 1995 | 2 |
| | 1 | NE Atlantic | Demersal trawl | 1988 | Northridge, 1988 | 3 |
| | 9+ | NE Atlantic | Trawl | 1971 – 1994 | Duguy, 1977; Duguy & Hussenot, 1982; Northridge 1988; Reijnders & | |
| | | | | | Lankester, 1990; Lowry, pers. com., 1994; Couperus, 1994; Nores et al., | |
| | | | | | 1992 in Fertl & Leatherwood, 1995 | |
| | 3 | French Atlantic coast | Bottom or mid- | 1971 – 1981 | Duguy & Hussenot, 1982 | |
| | | | water trawls | | | |
| | 2 | French Atlantic coast | Other fisheries | 1971 – 1981 | Duguy & Hussenot, 1982 | |
| | 1000+ | Faeroes | | | Northridge, 1984 | 4 |
| | Unknown | W Central Atlantic | | | NMFS, 1995 in Fertl & Leatherwood, 1995 | |
| | 1 | SW Atlantic | | | By Goodall et al., 1988, in Fertl & Leatherwood, 1995 | |
| | 1 | SW Pacific | trawl | | Mattlin pers. comm., 1995, in Fertl & Leatherwood 1995 | |
| White-beaked | A few | North Sea | trawl | Each year | Leatherwood & Reeves, 1983, in Northridge, 1984 | |
| dolphin | | | |] | | |
| • | 1 | North Sea | trawl | 1958 | Van Bree & Nijssen, 1964, in Fertl & Leatherwood, 1995 | |
| | unknown | NE England/North | Midwater herring | Every Summer | Northridge, 1988, in Northridge, 1991 | |
| | | Sea | trawls | | | |
| White-sided dolphin | 25 | NW Atlantic | Mackerel and | 1978 - 1991 | O'Hara et al., 1986; Young et al., 1993; NMFS, 1995, in Fertl & | |

¹ From the fleets of 8 nations fishing *in* the EEZ off the north-east United States coast.

² 142 animals *in* 1988, and 107 individuals *in* 1990 (according to Waring *et al.*, 1990; NMFS, 1995; Young *et al.*, 1993, Northridge, 1988).

³ Such an isolated account, as other ones related to other gears, are possibly indicative of sporadic accidental mortalities, but further information on the scale of these mortalities or their impact on populations is lacking (Northridge, 1991).

There has been a certain amount of exploitation of pilot whales, notably *in* the Faeroe Islands, where over 1000 are taken a year (Northridge, 1984).

| | | <u> </u> | squid trawls | | Leatherwood, 1995 | |
|-----------------------------|------------------------|--------------------|--------------------------------------|-----------------------|--|---|
| | 1 | NE England | Trawl or seine net | 1985 | IWC, 1987, in Northridge, 1991 | 5 |
| | No number | NE Atlantic | | | Couperus, 1994, in Fertl & Leatherwood, 1995 | |
| Dusky dolphin | 3 a few | SW Atlantic | Surface trawl | 1989 each year | Scialabba, 1989, in Northridge, 1991 and in Fertl & Leatherwood, 1995 | |
| | unknown | Chubut | trawl | Each year | Crespo & Corcuera, 1990, in Northridge, 1991 | |
| | 1 | Chubut | İ | 1989 | Crespo et al., 1994, in Fertl & Leatherwood, 1995 | İ |
| | Small numbers | SE Atlantic | trawl | | Northridge, 1984 | |
| | 11 | SE Atlantic | İ | 1989 - 1990 | Best pers. comm., 1992, in Fertl & Leatherwood, 1995 | İ |
| | occasional | SW Pacific | | | Slooten & Dawson, 1988, in Fertl & Leatherwood, 1995 | |
| | 3 | SW Pacific | | 1986 - 1988 | Anonymous 1988, in Fertl & Leatherwood, 1995 | |
| Peale's dolphin | No number | SW Atlantic | | | Crespo & Corcuera, 1990; Goodall et al., 1990, in Fertl & Leatherwood, 1995 | |
| Pacific White-sided dolphin | 3 | NE Pacific | | | Barlow et al., 1994, in Fertl & Leatherwood, 1995 | |
| | 11 | E Central Pacific | | 1973 - 1990 | Perez & Loughlin, 1991; NMFS, 1995, in Fertl & Leatherwood, 1995 | |
| Bottlenose dolphin | 3 | Mid Atlantic | Mackerel and squid fisheries | In eight years | Northridge, 1991 | |
| | 30 of which 17 in year | NW Atlantic | | 1977 – 1993 1993 | Waring et al., 1990; NMFS, 1995, in Fertl & Leatherwood, 1995 | |
| | 9+ | NE Atlantic | | 1971 – late 1980's | Mitchell, 1975; Duguy, 1977; Duguy & Hussenot, 1982; Anonymous, 1988; Currey <i>et al.</i> , 1990; Northridge, 1991, <i>in</i> Fertl & Leatherwood, 1995 | |
| | 6+ | W Central Atlantic | | 1976 - 1988 | Gunter, 1942; Prescott <i>et al.</i> , 1980; Burn & Sott, 1988; Reynolds, 1985; Gruber, 1981; Leatherwood & Reeves, 1982; Pharr, pers. com., 1991; Fertl, 1994; Wang <i>et al.</i> , 1994, <i>in</i> Fertl & Leatherwood, 1995 | |
| | Small numbers | Gulf of Mexico | Shrimp trawls, Gill nets and shot | | Northridge, 1991 | |
| Bottlenose dolphin Cont.d. | unknown | E Central Atlantic | Trawl, explosives | Pre 1957 | Cadanet, 1957, in Northridge, 1984 | 6 |
| | unknown | Mauritanian coast | Pelagic trawl | | Northridge, 1991; Scialabba, 1989, in Fertl & Leatherwood, 1995 | |
| | 4+ | Mediterranean | | | Duguy & Hussenot, 1982; Di Natale, 1989; Silvani et al., 1992, in Fertl & Leatherwood, 1995 | 7 |
| | Frequently | Adriatic part of | | | Mitchell, 1975, in Northridge, 1984 | 8 |

⁵ Catches are reported *in* the north-west Atlantic *in* foreign mackerel and squid trawl fisheries, but infrequently.

⁶ Cadanet (1957) refers to a troupe of bottlenose dolphins attacking a trawl net, which had to be frightened off with explosive charges (in Northridge, 1984).

⁷ According to Currey *et al.*, 1990, *in* Fertl and Leatherwood, 1995. incidental catches may be high *in* this area, although some dolphins are reported to be freed.

| | caught | Mediterranean | | | | |
|-------------------------|----------------|----------------------|-------------------|-------------|---|-----------------|
| | unknown | SW Atlantic | Midwater trawl | | Northridge, 1984 | |
| | | Argentinian coast | | | | |
| | 1 | | | | Goodall et al., 1988, in Fertl ad Leatherwood, 1995 | |
| | Small numbers | SE Atlantic | trawl | mid 1980 | Northridge, 1984 | |
| | | | | onwards | | |
| | 3 | W Indian Ocean (Gulf | | 1980 - 1982 | Pillai & Kasinathan, 1987, in Fertl & Leatherwood, 1995 | |
| | | of Mannar, India) | | | | |
| | 3 | E Indian Ocean | | Pre 1990 | Corkeron et al., 1990, Anonymous, 1991, in Fertl & Leatherwood, 1995 | |
| | including 1 in | (Australia) | | | | |
| | 1989 | | | | | |
| | Hundreds | SW Pacific | Midwater trawls | Each year | Mitchell, 1975, in Northridge, 1984 and in Fertl & Leatherwood, 1995 | |
| | suggested | | | | | |
| Risso's dolphin | A few | NW Atlantic | Trawl (mackerel | | O'Hara et al., 1986, in Northridge, 1991 | Such incidents |
| | 4 | NIXI A (1 (* | and squid) | 1005 1000 | W | seem to be rare |
| | 4 | NW Atlantic | | 1985 - 1992 | Waring et al., 1990 and NMFS, 1995, both in Fertl & Leatherwood, 1995 | |
| | A few | NW Atlantic | | | O'Hara et al., 1986, in Fertl & Leatherwood, 1995 | |
| | Occasional | NE Atlantic | | | Northridge, 1984; Reijnders & Lankester, 1990, in Fertl & Leatherwood, | |
| | 1 1 | W.C 1 Ad. d | | | 1995 | <u> </u> |
| | unnumbered | W Central Atlantic | | | NMFS, 1995, in Fertl & Leatherwood, 1995 | |
| Risso's dolphin cont.d. | 1 | SE Atlantic | | | Best pers. comm., 1992, in Fertl & Leatherwood, 1995 | |
| | 1 | NE Pacific | | 1973 - 1988 | Perez & Loughlin, 1991, in Fertl & Leatherwood, 1995 | |
| Spinner dolphin | Unknown, | SW Atlantic | | | Northridge, 1984 | |
| | probably few | (Argentina) | | | | |
| Dwarf spinner | 10 | W Central Pacific | | 1970/71 | Perrin et al., 1989, in Northridge, 1991 and in Fertl & Leatherwood, 1995 | 9 |
| dolphin | | (Gulf of Thailand) | | | | |
| Striped dolphin | 2 | NW Atlantic | | 1991 | NMFS 1995, in Fertl & Leatherwood, 1995 | |
| | 1 | NE Atlantic | | 1971 - 1976 | Duguy, 1977, in Fertl & Leatherwood, 1995 | |
| | unnumbered | NE Atlantic | | | Northridge, 1984, in Fertl & Leatherwood, 1995 | Occasionally |
| | | | | | | captured |
| | A large | Mediterranean | Trawl, tangle and | | Di Natale, 1989 in Northridge, 1991 | 10 |

⁸ One of the more frequently caught species *in* fishing nets, there has also been a live capture fishery *in* the Adriatic part of Mediterranean Sea (Mitchell 1975, *in* Northridge 1984) and a directed fishery *in* the Black Sea which was stopped *in* 1983. The Black Sea population of bottlenose dolphins may have been dramatically affected by the fishery (Smith 1976, *in* Northridge, 1984); there are no estimates of the current population size, but catches *in* recent years were probably *in* the tens of thousands, as since 1967 between 56,000 and 222,000 animals of 3 species (bottlenose dolphin, common dolphin and harbour porpoise) were being taken per annum (IWC 1983: Rep. of the Sub-Committee on small cetaceans)

⁹ A newly described dwarf form of this species from the Gulf of Thailand was identified on the basis of 10 specimens taken *in* shrimp trawls *in* the Gulf of Thailand *in* 1970/71.

| | number up to 5000 cetaceans | | longline | | | |
|----------------------------|--------------------------------------|-----------------------------------|---|---------------------------|--|----|
| | Unknown, probably few | SW Atlantic Argentina | Midwater trawl | | Northridge, 1984 | |
| Striped and common dolphin | 500 - 1000 | E Central Atlantic | | Per year | Maigret, 1994, in Fertl & Leatherwood, 1995 | |
| Common dolphin | 226 66, 76 | NW Atlantic | trawl | 1977 – 1992 1985, 1986 | O' Hara <i>et al.</i> , 1986; Waring <i>et al.</i> , 1990; NMFS, 1995, <i>in</i> Fertl & Leatherwood, 1995 | 11 |
| | 40 | NE Atlantic | | 1971 - 1982 | Duguy, 1977; Duguy & Hussenot, 1982; Pascoe, 1986; Nores <i>et al.</i> , 1992; Sequiera & Ferreira, 1994, <i>in</i> Fertl & Leatherwood, 1995 | |
| | Unknown | NE Atlantic | | | Reijnders & Lankester, 1990; Couperus, 1994, both <i>in</i> Fertl & Leatherwood, 1995 | |
| | 16 | French Atlantic coast | | 1981 | Collet, 1983, in Northridge, 1984 | |
| | Maybe hundreds | NE Atlantic | | | Northridge, 1988, <i>in</i> Northridge, 1991 Based on a very few number of interviews with fishermen | |
| | unknown | NE Atlantic | Midwater trawl and unidentified | | Northridge, 1988 and Mac Brearty, pers. comm., in Northridge, 1991 | |
| | unknown | Mauritania | Pelagic trawls | | Northridge, 1991 | |
| | Up to 6 – 22 per haul at night | E Central Atlantic, Mauritania | Surface sardine and mackerel trawls | | The Report of the Small Cetacean Sub-Committee, IWC 1982 pp113 – 126; in fishing Northridge, 1984; Maigret, 1994, in Fertl & Leatherwood, 1995 | 12 |
| | 1 | Mediterranean | | | Di Natale, 1983, in Fertl & Leatherwood, 1995 | |
| | unknown | Mediterranean | | 1989 | Di Natale, 1983, in Fertl & Leatherwood, 1995 | |
| | Extra statistics | Mediterranean | Sardine fishermen | | Tomilin, 1967, in Northridge, 1984 | 13 |
| | Unknown | SW Atlantic | | | Crespo & Corcuera, 1990, in Northridge, 1991 and Fertl & Leatherwood, 1995 | |
| | Small numbers | SE Atlantic | trawl | | Northridge, 1984 | |

¹⁰ On the basis of interviews with several hundred fishermen involved *in* this fishery, he estimated (pers. comm.) that total cetacean catches may have exceeded 5000 cetaceans *in* some years, suggesting a very large number of striped dolphins were taken. The impact on the population is unknown (Northridge, 1991)

About 60% of the total were taken *in* the foreign squid trawl fishery which has now been terminated. Catches of perhaps 20 per year continue *in* the mackerel fishery (Cranmore, 1988, *in* Northridge, 1991).

¹² Common dolphins are killed *in* surface trawling for sardines and mackerel *in* this region. One trawler was said to have caught 6 – 22 dolphins per haul at night, *in* an area where 12 trawlers were fishing.

¹³ Tomilin, 1967 *in* Northridge relates that common dolphin is also known to help Mediterranean sardine fishermen, who follow the dolphins which are responsible for frightening the fish to the surface. He continues that on the coast of Anatolia where "the population of dolphins totals about one million" they consume one million tonnes of fish *in* 100 days (Northridge, 1984)

| | 5 | SE Atlantic | | | Best, pers. comm., 1992, in Fertl & Leatherwood, 1995 | |
|--------------------------|---------------------------------|---|-------------------------------|---------------------|---|---|
| | 2 | | | | Cockcroft, per. Com., 1992, in Fertl & Leatherwood, 1995 | |
| | 9 | | Pair trawl | 1978 | | |
| | 2 | SW Pacific | Trawl | | | |
| | 11 | New Zealand | Pair trawl | | Reports to the IWC in Northridge, 1984 | |
| | 12 | | Trawl | 1979 | | |
| | 1 | | Bottom trawl | 1980 | | |
| Common dolphin cont.d. | 165+ | SW Pacific | | 1978 - 1995 | Fertl & Leatherwood, 1995; Mitchell 1975; Anonymous, 1981, 1982, 1988, 1989, 1990, 1991; Slooten & Dawson, 1988; Baird, 1995; Matlin, per. Com., 1995 | |
| Atlantic Spotted dolphin | 2 2 | W Central Atlantic | | 1985 1988 | Ford, pers. com., 1991, in Fertl & Leatherwood, 1995 | |
| Heaviside's dolphin | 2 | SE Atlantic | Stern trawler (Portuguese) | | IWC, 1979, in Northridge, 1984 | |
| | Small numbers 1 | | | 1977 – 1979 1990 | Fertl & Leatherwood, 1995; Anonymous, 1979; Northridge, 1984, Best, pers com., 1992 | |
| Hectors dolphin | unknown | SW Atlantic | Midwater trawl | | Northridge, 1984 | |
| | 10's | SW Atlantic Chubut, Tierra del Fuego, Peninsula Valdez | trawl | Per year | Scialabba, 1989, in Northridge, 1991; Crespo & Corcuera, 1990, in Northridge, 1991. | |
| | unknown | SW Atlantic, Chubut, Santa Cruz | Bottom & midwater trawls | | Northridge, 1991 | |
| | 2+ | SW Atlantic | | | Goodall et al., 1988, 1990, in Fertl & Leatherwood, 1995. | |
| Harbour porpoise | 1 | NW Atlantic | | 1977 1982 | Testaverde, 1978; O'Hara et al., 1986, in Fertl & Leatherwood, 1995 | |
| | Considerable nrs poss. hundreds | NE Atlantic | Danish trawls | annually | Anderson & Clausen, 1983, in Northridge, 1984 | |
| | 149 | Swedish waters | 86% in gillnets, | 1973 – 1987 | Lindstedt & Lindstedt, 1989, in Northridge, 1991 | |
| | 23 | | rest in trawls | 1988 | | |
| | >20 | Shetland | Bottom trawls | 1988/89 | Northridge, 1991 | |
| | unknown | British waters | trawl | | Northridge, 1991 | |
| | 1 | Baltic Sea | | 1950's | Fertl & Leatherwood, 1995; Anderson & Clausen, 1983; Anonymous, 1988, 1989, 1990; Lindstedt & Lindstedt, 1989; Kinze, 1994; Skora <i>et al.</i> , 1988; | - |
| | 244 | NE Atlantic | | 1973 - 1992 | Benke et al., 1991; Currey et al., 1990; Gaskin, 1984; Mitchell, 1975; | |

| | | | | | Moreno, 1993; Reijnders & Lankester, 1990; Northridge, 1988, 1991. | |
|---------------------|--------------|--------------------|-----------------|-------------|--|----|
| unknown | unknown | Black Sea | unknown | | Celikkale, 1990, in Northridge, 1991 | 14 |
| Harbour porpoise | infrequent | NE Pacific, Alaska | trawl | | Barlow et al., 1990 | 15 |
| | 1 | NE Pacific | Dredged by | 1941 | Fertl & Leatherwood, 1995; Scheffer & Slipp, 1948; Perez & Loughlin, | |
| | 8 | | trawler | 1973 - 1988 | 1991; Barlow et al., 1994 | |
| | unnumbered | E Central Pacific | | | Leatherwood & Reeves, 1986, in Fertl & Leatherwood, 1995 | |
| Vaquita | 6 | E Central Pacific | | 1961 – 1990 | Norris & Prescott, 1961; Vidal, 1990; Silber, pers. com., 1993, in Fertl & | |
| - | unnumbered | | | 1984 | Leatherwood, 1995; | |
| Spectacled porpoise | Unknown, | SW Atlantic | Midwater trawls | | Northridge, 1984 | |
| | probably few | Argentina | | | | |
| | unknown | Chubut and Santa | | | Crespo & Corcuera, 1990, in Northridge, 1991 | |
| | | Cruz, Argentina | | | | |
| Dall's porpoise | unknown | NE Pacific | | | Barlow, et al., 1990, in Northridge, 1991 | |
| | 55 | NE Pacific | | 1973 - 1993 | Perez & Loughlin, 1991; NMFS, 1995; Baird et al., 1991; Barlow et al., | |
| | | | | | 1994. | |
| | 14 | E Central Pacific | | 1973 - 1992 | Perez & Loughlin, 1991; NMFS, 1995, in Fertl & Leatherwood, 1995 | |
| Finless porpoise | 2 | E Indian Ocean | Shrimp trawl | | Niazi, 1990, in Northridge, 1991 and in Fertl & Leatherwood, 1995 | |
| | | (Pakistan) | | | | |

 ¹⁴ Trawls and purse seines are the major gear types *in* Turkish water
 15 As elsewhere *in* its range, this species appears to be widely affected by interactions with fisheries (Northridge, 1991).